

Conservation Evaluation of Stoloniferous Pussytoes, *Antennaria flagellaris*, in Canada*

GEORGE W. DOUGLAS¹, JENIFER L. PENNY², and KSENIA BARTON³

¹ Deceased.

² Conservation Data Centre, British Columbia Ministry of Environment, Ecosystems Branch, P.O. Box 9993, Stn Prov Govt, Victoria, British Columbia V8W 9R7 Canada

³Current address: 204-4272 Albert Street, Burnaby, British Columbia V5C 2E8 Canada

Douglas, George W., Jenifer L. Penny, and Ksenia Barton. 2006. Conservation evaluation of Stoloniferous Pussytoes, *Antennaria flagellaris*, in Canada. *Canadian Field-Naturalist* 120(2): 183–187.

In Canada, Stoloniferous Pussytoes, *Antennaria flagellaris*, is restricted to the Similkameen River valley south of Princeton, in south-western British Columbia. The three populations represent the northern limits of the species which ranges from south-western British Columbia, south in the western United States to Idaho, Wyoming, Nevada and California. In British Columbia, *Antennaria flagellaris* is associated with eroded, unstable, calcareous clay seepage slopes on open, southerly aspects. This habitat is infrequent in the Similkameen River area and the few existing plant populations could easily be extirpated through slight changes in drainage through drilling for coalbed methane gas, road-building, or housing development.

Key Words: Stoloniferous Pussytoes, *Antennaria flagellaris*, endangered, distribution, population size, British Columbia.